

# *FC* ELEMENTS OF ALGEBRAS AND ORDERS

C. Polcino Milies

## Abstract

Let  $\mathcal{U}(R)$  denote the group of units of an associative ring with unity  $R$ . We study elements of  $R$  which have a finite conjugacy class under the action of the elements of  $\mathcal{U}(R)$ . In particular, we give a survey of known results in the case when  $R$  is a group ring and state some new results for algebras and orders.

## Resumo

Seja  $\mathcal{U}(R)$  o grupo de unidades de um anel associativo com unidade  $R$ . Estudamos elementos de  $R$  que têm classe de conjugação finita sob a ação dos elementos de  $\mathcal{U}(R)$ . Em particular, descrevemos os resultados conhecidos no caso em que  $R$  é um anel de grupo e enunciamos alguns resultados novos para álgebras e ordens.

## 1 Introduction

Given an associative ring  $R$  with unity, we shall denote by  $\mathcal{U}(R)$  the group of units of  $R$ ; i.e., the set of invertible elements of  $R$ . We recall that an *FC group* is a group  $G$  such that all of its elements have finite conjugacy classes in  $G$ . More generally, we denote by  $\Phi(G)$  the *FC-center* of  $G$ , that is:

$$\Phi(G) = \{g \in G \mid [G : \mathcal{C}_G(g)] < \infty\}.$$

I. N. Herstein showed in [17] that if  $D$  is a division ring then  $\Phi(\mathcal{U}D)$  coincides with  $\mathcal{Z}(\mathcal{U}D)$ , the centre of  $\mathcal{U}D$ . The study of the FC-center of groups of units of group rings started with papers by S. K. Sehgal and H. J. Zassenhaus [26], C.

---

This work was partially supported by CNPq, Proc. 300243/79-0(RN).  
*1991 Mathematics Subject Classification:* Primary 16U60; Secondary 16H05, 20F24.  
*Key words and phrases:* algebras, orders, units, finite conjugacy.

Polcino Milies [21] and G. Cliff and S. K. Sehgal [8]. Also, A. Williamson [30], studied elements of a periodic group  $G$  which have finite conjugacy class in the group of units of its integral group ring. These results also follow from a paper by A. A. Bovdi [3]. A more general approach was given by S. K. Sehgal and H. J. Zassenhaus in [27]. This work was followed by several papers studying group rings over fields [22], [12].

In this short survey, we recall the origins of the theory of FC groups, then give a general view of the existing results about group rings with FC unit groups and, finally, state similar results in the more general context of algebras and orders.

## 2 FC groups

The results in this section are now standard. They are included here, with references to the original papers where they were published, only to serve as a historical introduction to the subject.

In 1948, R. Baer [1] introduced a series of finiteness conditions for groups which we list below:

- **(FC)** *Every element in the group  $G$  possesses only a finite number of conjugates in  $G$ .*
- **(LF)** *Every element in the group  $G$  is contained in a finite normal subgroup.*
- **(FO)** *There exists only a finite number of elements of any given order in the group  $G$ .*

Exploring the relations among these concepts, he obtained the following results.

**Theorem 2.1** *A group  $G$  is (LF) if and only if it is (FC) and contains no element of infinite order.*

As a consequence, it is easy to see that **(FO)** implies **(LF)** which, in turn, implies **(FC)**.

**Theorem 2.2** *A group  $G$  is FC if and only if every element is contained in a finitely generated normal subgroup and  $G/\mathcal{Z}(G)$  is LF, where  $\mathcal{Z}(G)$  denotes the centre of  $G$ .*

**Theorem 2.3** *A group  $G$  is FO if and only if  $\mathcal{Z}(G)$  is FO and, for every prime  $p$ , the factor group  $G/\mathcal{Z}(G)$  contains only a finite number of elements of order a power of  $p$ .*

In 1951, B. H. Neumann [19] continued the study of FC groups and gave their fundamental properties:

**Theorem 2.4** *Let  $G$  be an FC group. Then, the set  $T$  of elements of finite order in  $G$  is a characteristic subgroup of  $G$ ,  $G' \subset T$  and thus  $G/T$  is an abelian torsion-free group. Moreover, if  $G$  is finitely generated, then  $T$  is finite.*

This result has several consequences. Among these, we quote:

- If  $G$  is generated by a set of elements of finite order, then  $G$  is periodic (i.e.,  $G=T$ ). Moreover, if  $G$  is finitely generated, then  $G$  is finite.
- The torsion subgroup  $T$  is locally finite.
- (Baer)  $G$  is a periodic FC group if and only if  $G$  is LF.

**Theorem 2.5** *If  $[G : \mathcal{Z}(G)]$  is finite, then  $G'$  is finite. If  $G'$  is finite, then  $G$  is an FC group.*

FC centres of groups can be used to define a chain of subgroups and develop a theory similar to that of nilpotency. This was done in 1953 by F. Haimo [16]:

Set  $\Phi_1(G) = \{g \in G \mid [G : \mathcal{C}_G(g)] < \infty\}$ , the FC centre of  $G$  and, inductively, let  $\Phi_{n+1}(G)$  be the subgroup of  $G$  such that:

$$\frac{\Phi_{n+1}(G)}{\Phi_n(G)} = \Phi_1 \left( \frac{G}{\Phi_n(G)} \right).$$

The sequence of subgroups

$$\Phi_1(G) \subset \Phi_2(G) \subset \cdots \Phi_n(G) \subset \cdots$$

is called the *FC chain* of  $G$ . If there exists an integer  $n$  such that  $\Phi_n(G) = G$  and  $\Phi_{n-1}(G) \neq G$ , we say that  $G$  is *FC-nilpotent* of *FC-class* equal to  $n$ .

The following results show the similarity of these ideas with ordinary nilpotency:

**Theorem 2.6** *Let  $N$  be a normal subgroup of a group  $G$  such that  $N \subset \Phi_n(G)$ , for some integer  $n$  and such that there exists a positive integer  $k$  for which  $G/N$  is FC-nilpotent of class  $k$ . Then,  $G$  is FC-nilpotent of FC-class  $c \leq n + k$ .*

**Corollary 2.7** *If  $G' \subset \Phi_n(G)$  for some  $n$ , then  $G$  is FC-nilpotent of FC-class  $c \leq n + k$ .*

Shortly afterwards, J. Erdős [15] gave simpler proofs for some of Neumann's earlier results and also proved the following.

**Theorem 2.8** *If  $G$  is a finitely generated FC group, then  $[G : \mathcal{Z}(G)]$  is finite and thus  $G'$  is finite.*

He also used the theory of FC groups to prove a result of Ju. G. Fjodorov:

**Theorem 2.9** *If in an infinite group  $G$  every subgroup containing at least two elements is of finite index, then  $G$  is a cyclic group.*

Approximately at the same time, B. H. Neumann [20] studied *bounded* FC groups and proved the following.

**Theorem 2.10** *The number of elements in each conjugacy class of a group  $G$  is bounded if and only if the derived group  $G'$  is finite.*

The theory of FC groups has attracted considerable attention and is well developed. The interested reader may consult the book by M. J. Tomkinson [28] or the more recent survey [29].

### 3 FC elements in group rings

Given a ring  $R$  and a group  $G$ , we shall denote by  $RG$  the *group ring of  $G$  over  $R$* . The question of when the unit group  $\mathcal{U}(RG)$  is FC was first considered by S. K. Sehgal and H. J. Zassenhaus in 1977 [26] when they characterized groups  $G$  such that  $\mathcal{U}(\mathbb{Z}G)$  is FC:

**Theorem 3.1** *Let  $G$  be a group and let  $T$  denote the set of elements of finite order in  $G$ . Then  $\mathcal{U}(\mathbb{Z}G)$  is FC if and only if one of the following conditions holds:*

- (i)  $T$  is central in  $G$ .
- (ii)  $T$  is abelian, non-central and, for all  $t \in T$  and all  $x \in G$  we have that  $xtx^{-1} = t^{\pm 1}$ .
- (iii)  $T = E \times \mathcal{Q}_8$ , where  $E$  is an elementary abelian 2-group,

$$\mathcal{Q}_8 = \langle a, b \mid a^4 = 1, a^2 = b^2, aba^{-1} = b^{-1} \rangle$$

and conjugation by any element  $x \in G$  induces an inner automorphism of  $\mathcal{Q}_8$ .

With this result, they were able to characterize also FC unit groups of  $KG$ , when  $K$  is a field of characteristic 0. The case when  $\text{char}(K) = p > 0$  was completed in a sequence of two papers, by C. Polcino Milies [21] and G. Cliff and S. K. Sehgal [8]. The characterizations involved the condition of every idempotent of  $KT$  being central in  $KG$ . After this condition was better understood in [9], [10] and [11], these results can be stated as follows.

**Theorem 3.2** *Let  $G$  be a torsion group. Then  $\mathcal{U}(KG)$  is an FC group if and only if either  $KG$  is finite or  $G$  is abelian.*

**Theorem 3.3** *Assume that  $\text{char}(K) = p > 0$  and that  $G$  is a non-torsion group which contains  $p$ -element. Then  $\mathcal{U}(KG)$  is an FC group if and only if either  $G$  is abelian or  $G$  is a non-abelian FC group,  $p = 2$ ,  $T = \langle t \rangle \times A$  where  $o(t) = 2$ ,  $A$  is a finite group of odd order,  $G' = \langle t \rangle$  and  $T$  is central.*

For a given field  $K$  we shall denote by  $P(K)$  the prime subfield of  $K$ .

**Theorem 3.4** *Assume that  $\text{char}(K) = p > 0$  and that  $G$  is a non-torsion group which contains no  $p$ -elements. Then  $\mathcal{U}(KG)$  is an FC group if and only if either  $G$  is abelian or  $G$  is a non-abelian FC group,  $T$  is abelian and one of the following conditions holds:*

- (i)  *$KT$  is finite and for all  $t \in T$  and all  $x \in G$  we have that  $t^x = t^{p^r}$  for some non-negative integer  $r = r(x, t)$  which is a multiple of  $[K : P(K)]$ .*
- (ii)  *$T$  is finite, central.*
- (iii)  *$T$  is central, of the form  $T = \mathbb{Z}(q^\infty) \times B$  for some prime  $q \neq p$ ,  $G' \subset \mathbb{Z}(q^\infty)$  and there exists an integer  $k$  such that  $K$  does not contain roots of unity of order  $q^k$ .*

**Theorem 3.5** *Let  $\text{char}(K) = 0$  and assume that  $G$  is a non-torsion group. Then  $\mathcal{U}(KG)$  is an FC group if and only if  $G$  is either abelian or a non-abelian FC group with  $T$  central and, if  $T$  is infinite, then  $T$  and  $K$  can be described as in part (iii) of the previous theorem.*

Results on the construction of the group of units of a group ring, in general, can be found in [5]. The description of group algebras having FC unit groups was also given independently by A. A. Bovdi in [4]. See also [6].

## 4 The supercentre of a group

In 1978, A. Williamsom [30] studied elements of a group  $G$  which have a finite conjugacy class in the unit group of the integral group ring of  $G$ . He proved the following.

**Theorem 4.1** *Let  $G$  be a periodic group. An element  $x \in G$  has a finite conjugacy class in  $\mathcal{U}(\mathbb{Z}G)$  if and only if either:*

- (i)  *$x$  is central in  $G$ , or*

(ii)  $o(x) = 4$  and  $x$  belongs to an abelian group  $H$  of index 2 in  $G$ , with

$$G = \langle H, c \mid c^2 = x^2, h^c = h^{-1}, \forall h \in H \rangle.$$

In 1981, S. K. Sehgal and H. J. Zassenhaus [27] defined the *FC subring* of a ring  $R$  as:

$$FC(R) = \{x \in R \mid \text{the conjugacy class of } x \text{ under } \mathcal{U}(R) \text{ is finite}\}.$$

In that same paper, they also introduced the *supercentre* of a group  $G$  as:

$$\begin{aligned} S(G) &= G \cap FC(\mathbb{Z}G) = G \cap \Phi\mathcal{U}(\mathbb{Z}G) \\ &= \{g \in G \mid \text{the conjugacy class of } g \text{ in } \mathcal{U}(\mathbb{Z}G) \text{ is finite}\}. \end{aligned}$$

**Theorem 4.2** *Let  $G$  be a finite group. Then, the FC subring of  $\mathbb{Z}G$  consists of all those elements  $x \in \mathbb{Z}G$  such that, for every irreducible representation  $f$  of  $\mathbb{Q}G$  over  $\mathbb{Q}$  for which  $f(\mathbb{Q}G)$  is not a totally definite quaternion algebra, we have that  $f(x)$  is central in  $f(\mathbb{Q}G)$ .*

**Theorem 4.3** *Let  $G$  be a finite group. Then;*

$$T(\Phi(\mathcal{U}(\mathbb{Z}G))) = \pm S(G).$$

This last theorem is also contained in a paper by A. A. Bovdi [2].

Shortly afterwards, C. Polcino Milies and S. K. Sehgal [22] defined, for a group  $G$  and an arbitrary ring  $K$ , the  *$K$ -supercentre* of  $G$  as:

$$S_K(G) = G \cap FC(KG) = G \cap \Phi(\mathcal{U}(KG)).$$

A complete description of supercentres of groups over fields was given by S. P. Coelho and C. Polcino Milies [12].

## 5 FC centres of algebras and orders

Let  $D$  be an integral domain,  $K$  its field of fractions and  $A$  an algebra over  $K$ . We recall that an unital subring  $\Lambda$  of  $A$  is said to be a  *$D$ -order* in  $A$  if it

has a finite basis when considered as a  $D$ -module and  $K\Lambda = A$ . The results that follow, concerning the FC-subring of algebras and orders were announced in [13] and the proofs will appear in [14].

The key remark for the results on FC-subrings of algebras that will be given below is the following.

**Theorem 5.1** *Let  $G$  be a connected algebraic group. Then, every element with a finite conjugacy class is central.*

As a consequence, we obtain.

**Theorem 5.2** *Let  $A$  be an algebra with unity over an infinite field  $K$ .*

(i) *If  $A$  is finite dimensional, then  $\mathcal{U}A$  is a connected linear algebraic group and, consequently,*

$$\Phi(\mathcal{U}A) = \mathcal{Z}(\mathcal{U}A).$$

*Moreover,  $A$  is generated by its units, as a vector space over  $K$  and, therefore,  $\mathcal{U}A$  is FC if and only if  $A$  is commutative.*

(ii) *Every torsion unit of  $\Phi(\mathcal{U}A)$  commutes with each algebraic unit of  $A$  and, consequently,  $\Phi(\mathcal{U}A)$  is solvable of length at most 2.*

(iii) *Every element of  $\Phi(\mathcal{U}A)$  commutes with each nilpotent element of  $A$ .*

The results above were obtained using the fact that the group of units of a finite dimensional algebra can be viewed as a connected algebraic group. A more general version of these results, from another point of view, appears in [7].

For an order  $\Lambda$  in an algebra  $A$ , we have the following.

**Theorem 5.3** *Let  $D$  be an infinite domain,  $K$  its field of fractions,  $A$  a finite dimensional  $K$ -algebra,  $\Lambda$  a  $D$ -order in  $A$ ,  $\mathcal{J} = \mathcal{J}(A)$  the Jacobson Radical of  $A$  and  $\overline{A} = A/\mathcal{J}$ . Assume that  $\text{Hom}_A(P_i, P_j) = 0$  for every pair of non-isomorphic principal modules  $P_i, P_j$  of multiplicity 1 in  $A$ . If every minimal ideal of  $\overline{A}$  which is a division ring is isomorphic to  $K$ , then*

$$\Phi(\mathcal{U}\Lambda) \subset \mathcal{Z}(A),$$



**Corollary 5.4** *Let  $D$  and  $K$  be as above,  $A$  be a finite dimensional  $K$ -algebra and  $\Lambda$  an order in  $A$ . Assume that  $\text{Hom}_A(P_i, P_j) = 0$  for every pair of non-isomorphic principal modules  $P_i, P_j$  of multiplicity 1 in  $A$ . If  $K$  is a splitting field for  $A$ , then*

$$\Phi(\mathcal{U}\Lambda) \subset \mathcal{Z}(A).$$

**Corollary 5.5** *Let  $D$  and  $K$  be as above,  $A$  be a semisimple finite dimensional  $K$ -algebra and  $\Lambda$  a  $D$ -order in  $A$ . If  $A$  has no minimal ideal which is a non-commutative division ring then*

$$\Phi(\mathcal{U}\Lambda) \subset \mathcal{Z}(A).$$

**Theorem 5.6** *Let  $D$  be an infinite domain and  $R$  a  $D$ -algebra.*

(i) *If  $R$  is torsion free as a  $D$ -module then*

$$\Phi(GL_n(R)) = \Phi(\mathcal{U}R)I,$$

where  $I$  is the identity matrix of  $M_n(R)$ .

(ii) *If  $\text{char}(D) = 0$  and  $n > 1$  then*

$$\Phi(GL_n(R)) = \mathcal{Z}(GL_n(R)).$$

Applying this results in the context of group rings, we may obtain the following.

**Lemma 5.7** *Let  $K$  be a field and let  $G$  be a subgroup of  $GL(2, K)$ . Then*

(i) *if  $a \in GL(2, K)$  is noncentral, its centralizer in  $GL(2, K)$  is abelian and*

(ii) *either  $\Phi(G) = \mathcal{Z}(G)$  or  $G$  is abelian-by-finite.*

**Proposition 5.8** *Let  $G = K_8 \times \langle c \rangle$ , where  $c$  is an element of order  $p$ , an odd prime, and  $K_8 = \langle a, b \rangle$  is the quaternion group of order 8. Then*

$$\Phi(\mathcal{U}(\mathbb{Z}G)) = \mathcal{Z}(\mathcal{U}(\mathbb{Z}G)).$$

**Proposition 5.9** *Let  $G$  be a finite group and assume that  $T\Phi(\mathcal{U}(\mathbb{Z}G))$  is non-abelian. Then  $G$  is a 2-group.*

The theorem of Williamson quoted above follows easily from these results.

## References

- [1] Baer, R., *Finiteness properties of groups*, Duke J. of Math., 15 (1948), 1021-1032.
- [2] Bovdi, A. A., *The periodic normal divisors of the multiplicative group of a group ring*, Sibirski Mat. Z., 9, 2 (1968), 495-498.
- [3] Bovdi, A. A., *On the conjugacy classes of an integral group ring*, Canad. Math. Bull., 21 (1978), 491-496.
- [4] Bovdi, A. A., *Construction of the multiplicative group of a group algebra with finiteness conditions*, Mat. Issled, 56 (1980), 14-27.
- [5] Bovdi, A. A., *The structure of the multiplicative group of an integral group ring*, Dokl. Acad. Nauk SSSR, 301 (1988), 1295-1297.
- [6] Bovdi, V., *Twisted group rings whose units form an FC group*, Canad. J. Math, 47 (1995), 274-289.
- [7] Bovdi, V., *On elements in algebras having finite number of conjugates*, Publ. Math. Debrecen, 57 (2000), 231-239.
- [8] Cliff, G. H.; Sehgal, S. K., *Group Rings whose Units form an FC group*, Math. Z., 161 (1978), 163-168.
- [9] Coelho, S. P., *A note on central idempotents in group rings*, Proc. Edinburgh Math. Soc., 30 (1987), 60-72.
- [10] Coelho, S. P.; Polcino Milies, C., *A note on central idempotents in group rings II*, Proc. Edinburgh Math. Soc., 31 (1988), 211-215.

- [11] Coelho, S. P.; Polcino Milies, C., *Some remarks on central idempotents in group rings*, Publ. Math (Debrecen), 52 (1998), 187-192.
- [12] Coelho, S. P.; Polcino Milies, C., *Finite Conjugacy in Group Rings*, Comm. in Algebra, 19, 3 (1991), 981-995.
- [13] Dokuchaev, M. A.; Juriaans, S. O.; Polcino Milies, C.; Sobral Singer, M. L., *FC centers of units in algebras and orders*, C.R. Math. Rep. Acad. Sci. Canada, 22 (2000), 25-27.
- [14] Dokuchaev, M. A.; Juriaans, S. O.; Polcino Milies, C.; Sobral Singer, M. L., *Finite Conjugacy in Algebras and Orders*, Proc. Edinburgh Math. Soc., to appear.
- [15] Erdős, J., *The theory of groups with finite classes of conjugate elements*, Acta Math. Acad. Sci, Hungar., 5 (1954), 45-58.
- [16] Haimo, F., *The FC chain of a group*, Canad. J. of Math., 5 (1953), 498-511.
- [17] Herstein, I. N., *Conjugates in Division Rings*, Proc. A.M.S., 7 (1956), 1021-1022.
- [18] Herstein, I. N., *Finite multiplicative subgroups in division rings*, Pacific J. of Math., 1 (1953), 121-126.
- [19] Neumann, B. H., *Groups with finite classes of conjugate elements*, Proc. London Math. Soc., 1 (1951), 178-187.
- [20] Neumann, B. H., *Groups covered by permutable subsets*, J. London Math. Soc., 29 (1954), 236-248.
- [21] Polcino Milies, C., *Group Rings whose Units form an FC Group*, Archiv der Math., 30 (1978), 380-384.
- [22] Polcino Milies, C.; Sehgal, S. K., *FC elements in group rings*, Comm. in Algebra, 9 (1981), 1285-1293.

- [23] Ritter, J.; Sehgal, S. K., *Generators and Subgroups of  $\mathcal{U}(\mathbb{Z}G)$* , Contemporary Math., 93 (1989), 331-347.
- [24] Sehgal, S. K., *Topics in group rings*, Marcel Dekker, New York, 1978.
- [25] Sehgal, S. K., *Units in integral group rings*, Longman Scientific and Technical, Essex, 1993.
- [26] Sehgal, S. K.; Zassenhaus, H. J., *Group Rings whose Units form an FC Group*, Math. Z., 153 (1977), 29-35.
- [27] Sehgal, S. K.; Zassenhaus, H. J., *On the Supercentre of a Group and its Ring Theoretic Generalization*, in Integral Representations and its Applications, Lecture Notes in Mathematics No 882, Springer-Verlag, Berlin, 1981, pp. 117-144.
- [28] Tomkinson, M. J., *FC groups*, Pitman, Boston, 1984.
- [29] Tomkinson, M. J., *FC groups: recent progress*, in Infinite groups 1994 (Ravello 1994), Walter de Gruyter, Berlin, 1995, pp. 271-285.
- [30] Williamson, A., *On the conjugacy classes in an integral group ring*, Canad. Math. Bull., 21, 4 (1978), 491-496.

Instituto de Matemática e Estatística

Universidade de São Paulo

Caixa Postal 66281

05315-970 - São Paulo - Brasil

*E-mail: polcino@ime.usp.br*