

# A Variational Principle for the Specific Entropy for Symbolic Systems with Uncountable Alphabets

D. Aguiar<sup>§</sup>, L. Cioletti<sup>§</sup> and R. Ruviano<sup>§</sup>

<sup>§</sup>Universidade de Brasília, MAT, 70910-900, Brasília, Brazil

## Abstract

In this paper we derived a variational principle for the specific entropy on the context of symbolic dynamics of compact metric space alphabets and use this result to obtain the uniqueness of the equilibrium states associated to a Walters potential.

## 1 Introduction

Let  $\mathbb{N}$  be the set of positive integers,  $\mathcal{A}$  a finite alphabet,  $\Omega \equiv \mathcal{A}^{\mathbb{N}}$  the product space equipped with its usual distance  $d_{\Omega}$ , product topology and the sigma-algebra generated by the open sets. The dynamics in this paper is given by  $\sigma : \Omega \rightarrow \Omega$ , the left shift mapping. In this setting an *equilibrium state* for a continuous potential  $f : \Omega \rightarrow \mathbb{R}$  is an element of  $\mathcal{M}_{\sigma}(\Omega)$ , the set of all shift-invariant Borel probability measures, solving the following variational problem

$$\sup_{\mu \in \mathcal{M}_{\sigma}(\Omega)} \{H(\mu) + \int_{\Omega} f d\mu\}, \quad (1)$$

where  $H(\mu)$  is the Kolmogorov-Sinai entropy of  $\mu$ . This variational principle was introduced by Ruelle [Rue67] in the context of Statistical Mechanics and later Walters [Wal75] considered this problem in the Ergodic Theory setting. This classical problem still is a central one in Ergodic Theory/Thermodynamic Formalism and the complete classification of all continuous potentials for which the problem (1) has a unique solution remains open.

One of the goals of this paper is to establish the uniqueness of the equilibrium states for Hölder and Walters potentials (see Definition 5 for the former) in a more general setting, where the alphabet  $\mathcal{A}$  is a general compact metric space, thus including cases where  $\mathcal{A}$  is uncountable. We shall remark that this uniqueness result is well-known in the context of finite alphabets, see [Wal01, Bal00, PP90] and references therein. The

---

2010 *Mathematics Subject Classification*: 37D35.

*Keywords*: Thermodynamic Formalism, Ruelle operator, entropy, equilibrium states.

first step towards this uniqueness result on general compact metric space alphabets was given in [LMMS15]. They proved that any probability measure  $\mu$  satisfying  $\mathcal{L}_f^* \mu = \mu$  (see Section 3) is a solution to the variational problem and also that the set of such probability measures is a singleton. In the finite-alphabet case this implies, by a result in [PP90], the uniqueness of the equilibrium states. But it is not evident that the results from [PP90] can be applied in cases where  $\mathcal{A}$  is a general compact metric space. The main idea presented here to solve this problem is to use the Ruelle operator and DLR-Gibbs measures theory, suitably adapted for the one-dimensional one-sided lattice.

In what follows we recall how the Ruelle operator associated to a Hölder potential and the equilibrium states are linked, in the finite-alphabet context. Next we move the discussion to more general settings.

For each  $0 < \alpha < 1$ , we denote by  $C^\alpha(\Omega) \equiv C^\alpha(\Omega, \mathbb{R})$  the space of all real valued  $\alpha$ -Hölder continuous functions which is defined as usual by

$$\left\{ f : \Omega \rightarrow \mathbb{R} : \text{Hol}_\alpha(f) \equiv \sup_{x \neq y} \frac{|f(x) - f(y)|}{(d_\Omega(x, y))^\alpha} < \infty \right\}.$$

When the potential  $f$  has nice regularity properties such as  $\alpha$ -Hölder continuity, equilibrium states and fine properties of them can be obtained by using the Ruelle transfer operator  $\mathcal{L}_f$ , which is defined for each continuous function  $\varphi$  by the following expression

$$\mathcal{L}_f(\varphi)(x) = \sum_{a \in \mathcal{A}} \exp(f(ax)) \varphi(ax), \quad \text{where } ax \equiv (a, x_1, x_2, \dots).$$

If  $f$  is an  $\alpha$ -Hölder continuous function then the Ruelle-Perron-Frobenius theorem ensures, among other things, that  $\lambda_f$ , the spectral radius of  $\mathcal{L}_f$  acting on  $C^\alpha(\Omega)$ , is a positive maximal isolated eigenvalue and associated to it we have a strictly positive eigenfunction  $h_f$ . Moreover, there is a Borel probability measure  $\nu_f$  on  $\Omega$ , such that  $\mathcal{L}_f^* \nu_f = \lambda_f \nu_f$ , where  $\mathcal{L}_f^*$  is the Banach transpose of  $\mathcal{L}_f : C(\Omega) \rightarrow C(\Omega)$ , and  $C(\Omega) \equiv C(\Omega, \mathbb{R})$  is the space of all real continuous function from  $\Omega$ . In this case there is a unique equilibrium state for  $f$  and it is given, up to a normalization, by the probability measure  $\mu_f \equiv h_f \nu_f$ , see [Bal00, PP90, Rue68]. Besides solving the variational problem the spectral data of the Ruelle operator can also be used to obtain a variational formulation of the Kolmogorov-Sinai entropy of  $\mu_f$  as follows

$$H(\mu_f) = \log |\mathcal{A}| + \inf_{g \in C^\alpha(\Omega)} \left\{ - \int_\Omega g \, d\mu_f + \log \lambda_g \right\},$$

see [LMMS15]. If  $f \equiv 0$ , then  $H(\mu_f) = \log |\mathcal{A}|$  and therefore when the number of symbols in our alphabet goes to infinity, i.e.,  $|\mathcal{A}| \rightarrow \infty$ , the Kolmogorov-Sinai entropy of  $\mu_f$  will be infinity.

To handle the case of infinite alphabets, a new definition for entropy of a Gibbs measure associated to a Hölder potential is proposed in [LMMS15]. To circumvent the pointed-out problem with Kolmogorov-Sinai entropy, the authors considered an *a priori* Borel probability measure  $p$  on  $\mathcal{A}$  having full support, and next a  $p$ -dependent concept

of entropy is given. It takes non-positive values and attain its supremum in the product measure  $\prod_{i \in \mathbb{N}} dp$ . The introduction of an a priori measure is also a standard procedure, in Equilibrium Statistical Mechanics, when dealing with continuous spin systems, see [vEFS93, Geo11]. However, in [LMMS15] no mention to a possible relation between these two approaches is made. In [GKLM18] the authors develop an abstract theory of Thermodynamic Formalism, where entropy is defined as a kind of Legendre-Fenchel dual of the topological pressure. It worth to mention that similar results are well-known in the context of Statistical Mechanics, see for example [Isr79]. Here we show that the specific entropy coincides with the one introduced in [LMMS15] and it can be extended to the classical Legendre-Fenchel transform of the topological pressure, thus providing a concrete representation (as a Thermodynamic Limit) of this function in cases where the alphabet  $\mathcal{A}$  is a general compact metric space. Furthermore, we obtain (Theorem 1) a version of the important identity (15.32) of [Geo11]

$$\lim_{n \rightarrow \infty} \frac{1}{|\Lambda_n|} \mathcal{H}_{\Lambda_n}(\mu|\nu) = P(\Phi) + \langle \mu, P \rangle - \mathfrak{h}^s(\mu)$$

in the context of symbolic dynamics for potentials in the Walters class, see Definition 7. We also show that the entropy defined in [LMMS15] is equal to the specific entropy (or mean entropy or entropy rate) commonly used in Statistical Mechanics (see [Geo11]) and as by-product a variational formulation for the specific entropy is derived. Afterwards, these results are applied to prove uniqueness of equilibrium states for potentials in Walters class, when  $\mathcal{A}$  is a general compact metric alphabet.

We should mention that our results about the uniqueness of the equilibrium states for potentials in Walters space can not be deduced from the recent results in [GKLM18]. The first reason is related to the finite-to-one map hypothesis imposed there, which is broken when the alphabet is infinite. The second reason is that the Ruelle operator associated to a Walters potential does not have, in general, the spectral gap property, see [CS16]. In such cases the arguments given in [GKLM18] can not be used. Although the references [CL16, CS16, LMMS15] already considered compact alphabets and equilibrium states in such context, the uniqueness results discussed in these papers concern to the eigenmeasures of the Banach transpose of the Ruelle operator. As mentioned earlier, in the finite-alphabet case, the uniqueness of the eigenmeasures implies the uniqueness of the equilibrium states for Hölder and Walters potentials. For general compact metric alphabets more work is required. The additional results needed to prove the uniqueness of the equilibrium states are obtained in Section 3.

In [CL16, CS16, LMMS15] a concept of entropy is considered in the context of compact metric alphabets. It is defined by a variational principle, but it was not recognized as the negative of the Legendre-Fenchel transform of the pressure functional. Here we obtain as a byproduct of the proof of Theorem 3: the entropy considered in these works is actually the negative of the Legendre-Fenchel dual of the pressure functional and, moreover, is equal to the specific entropy. This equality, the content of Theorem 3, has a further consequence due to Proposition 15.14 in [Geo11] which reads: the entropy considered in these works is an affine function, when restricted to the subspace of shift-

invariant Borel probability measures. Although this was a known result for dynamical systems with finite topological entropy, it is new for uncountable alphabets.

## 2 Specific Entropy

In this section we recall the definition and basic properties of the specific entropy, the exposition follows closely the reference [Geo11].

From now on  $\mathcal{A}$  is a compact metric space and  $\mathcal{F}$  denotes the sigma-algebra of  $\Omega$  generated by the open sets. Let  $\mu, \nu$  be two probability measures on the measurable space  $(\Omega, \mathcal{F})$ . The negative of the *relative entropy* of  $\mu$  with respect to  $\nu$  on the sub-sigma-algebra  $\mathcal{A}$ , notation  $\mathcal{H}_{\mathcal{A}}(\mu|\nu)$ , is the extended real number

$$\mathcal{H}_{\mathcal{A}}(\mu|\nu) \equiv \begin{cases} \int_{\Omega} \frac{d\mu|_{\mathcal{A}}}{d\nu|_{\mathcal{A}}} \log \left( \frac{d\mu|_{\mathcal{A}}}{d\nu|_{\mathcal{A}}} \right) d\nu, & \text{if } \mu \ll \nu \text{ on } \mathcal{A}; \\ \infty, & \text{otherwise.} \end{cases}$$

For each finite  $\Lambda \subset \mathbb{N}$  consider the projection  $\pi_{\Lambda} : \Omega \rightarrow \mathcal{A}^{\Lambda}$  given by  $\pi_{\Lambda}(x) = (x_i)_{i \in \Lambda}$ . We denote by  $\mathcal{F}_{\Lambda}$  the sigma-algebra generated by the projections  $\{\pi_{\Gamma} : \Gamma \subset \Lambda\}$  and we define the relative entropy of  $\mu$  with respect to  $\nu$  in  $\mathcal{F}_{\Lambda}$  by  $\mathcal{H}_{\Lambda}(\mu|\nu) \equiv \mathcal{H}_{\mathcal{F}_{\Lambda}}(\mu|\nu)$ .

Fix an a priori probability measure  $p$  on  $\mathcal{A}$ . For each probability measure  $\mu$  on  $\Omega$  and a finite volume  $\Lambda \subset \mathbb{N}$ , the relative entropy of  $\mu$  in  $\Lambda$ , with respect to  $p$ , is defined by  $\mathcal{H}_{\Lambda}(\mu) \equiv -\mathcal{H}_{\Lambda}(\mu | \prod_{i \in \mathbb{N}} p)$ . If  $\Lambda_n \equiv \{1, \dots, n\}$  and  $\mu \in \mathcal{M}_{\sigma}(\Omega)$ , is shown in [Geo11] that the following limit

$$\mathbf{h}^s(\mu) = \lim_{n \rightarrow \infty} \frac{\mathcal{H}_{\Lambda_n}(\mu)}{n}$$

always exists in  $[-\infty, 0]$  and  $\mathbf{h}^s(\mu)$  is called the **specific entropy** per site of  $\mu$  relative to the a priori probability measure  $p$ . The specific entropy  $\mathbf{h}^s$  is always a concave and upper semicontinuous function. Since  $\mathbf{h}^s(\prod_{i \in \mathbb{N}} p) = 0$ , for any choice  $p$ ,  $\mathbf{h}^s$  is not identically constant equal to  $-\infty$ . An extreme case occurs when  $p$  is a Dirac measure concentrated on an arbitrary point of  $\mathcal{A}$ . In this case the product measure  $\prod_{i \in \mathbb{N}} p$  is the only  $\sigma$ -invariant probability measure for which this function take finite values. For the proof of these properties and more details on specific entropy, see [Geo11].

## 3 Main Results

In this section we obtain a variational formulation for the specific entropy and also prove the uniqueness of equilibrium states for a large class of potentials. For the sake of simplicity, we present the argument for Hölder potentials and point out, in last section, what are the needed changes to prove the theorem in more general cases.

Let  $(\mathcal{A}, d_{\mathcal{A}})$  be an arbitrary compact metric space and consider the symbolic space  $\Omega = \mathcal{A}^{\mathbb{N}}$  equipped with a metric  $d_{\Omega}$  which induces the product topology. For example,  $d_{\Omega}(x, y) \equiv \sum_{n=1}^{\infty} 2^{-n} d_{\mathcal{A}}(x_n, y_n) / (1 + d_{\mathcal{A}}(x_n, y_n))$ .

Given an a priori measure  $p$  on  $\mathcal{A}$  and an  $\alpha$ -Hölder continuous potential  $f$  we define the Ruelle operator  $\mathcal{L}_f : C^\alpha(\Omega) \rightarrow C^\alpha(\Omega)$  as being the linear operator sending  $\varphi$  to  $\mathcal{L}_f(\varphi)$ , which is given by the following expression

$$\mathcal{L}_f(\varphi)(x) = \int_{\mathcal{A}} \exp(f(ax))\varphi(ax) dp(a), \quad \text{where } ax \equiv (a, x_1, x_2, \dots).$$

A potential  $f$  is said to be normalized if  $\mathcal{L}_f(1)(x) = 1$  for all  $x \in \Omega$ . By using the generalization of Ruelle-Perron-Frobenius theorem provided in [LMMS15] we can associate to any Hölder potential  $f$  a cohomologous normalized potential  $\bar{f}$  given by

$$\bar{f} = f + \log h_f - \log(h_f \circ \sigma) - \log \lambda_f, \quad (2)$$

where  $\lambda_f$  is a maximal eigenvalue of  $\mathcal{L}_f$  and  $h_f$  is a strictly positive  $\alpha$ -Hölder eigenfunction associated  $\lambda_f$ . The authors also proved that  $\mathcal{G}^*(\bar{f}) \equiv \{\nu \in \mathcal{M}_1(\Omega) : \mathcal{L}_{\bar{f}}^* \nu = \nu\}$  is a singleton and contained in  $\mathcal{M}_\sigma(\Omega)$ . This measure is called here the Gibbs measure associated to the potential  $f$ .

Now we consider the following family of probability kernels  $(\gamma_n)_{n \geq 1}$ , where for each  $n \geq 1$  the kernel  $\gamma_n : \mathcal{F} \times \Omega \rightarrow [0, 1]$  is given by  $\gamma_n(A|y) \equiv \mathcal{L}_f^n(1_A)(\sigma^n(y))$ , where  $f \in C^\alpha(\Omega)$  is a normalized potential. For any probability measure  $\nu$ , on  $\Omega$ , and  $y \in \Omega$  we define a probability measure  $\nu\gamma_n(\cdot|y) \equiv \int_{\Omega} \gamma_n(\cdot|y) d\nu(y)$ . We remark that if  $\nu$  is such that  $\mathcal{L}_f^* \nu = \nu$  then for any  $A \in \mathcal{F}$  and  $y \in \Omega$  we have

$$\begin{aligned} \nu\gamma_n(A|y) &= \int_{\Omega} \mathcal{L}_f^n(1_A) \circ \sigma^n(y) d\nu(y) = \int_{\Omega} \mathcal{L}_f^n(1_A)(y) d\nu(y) \\ &= \int_{\Omega} 1_A(y) d[(\mathcal{L}_f^*)^n \nu](y) = \nu(A). \end{aligned}$$

Our next theorem is inspired by Theorem 15.30 of [Geo11]. The hypotheses there are not fully satisfied in our setting. Firstly, here we are working on one-dimensional one-sided lattice. Secondly, the family of kernels considered here is not defined by a uniformly summable translation invariant interaction. The main difference of our proof is the use of duality properties of the Ruelle operator.

**Theorem 1.** *Let  $p$  be an a priori probability measure on  $\mathcal{A}$  having full support and  $f \in C^\alpha(\Omega)$ . Then for each  $\mu \in \mathcal{M}_\sigma(\Omega)$  and  $\nu \in \mathcal{G}^*(\bar{f})$ , the following limit exists*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathcal{H}_{\Lambda_n}(\mu|\nu) \equiv \mathbf{h}(\mu|\nu) = \log \lambda_f - \int_{\Omega} f d\mu - \mathbf{h}^s(\mu).$$

*Proof.* If for some  $n \in \mathbb{N}$ , we have  $\mathcal{H}_{\Lambda_n}(\mu|\nu) = +\infty$ , then  $\mathcal{H}_{\Lambda_n}(\mu|\nu\gamma_n(\cdot|y)) = +\infty$  and therefore  $\mathcal{H}_{\Lambda_n}(\mu|\gamma_n(\cdot|y)) = +\infty$ , so  $\mu|_{\mathcal{F}_{\Lambda_n}}$  is not absolutely continuous with respect to  $\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}$ . Since  $f$  is bounded, it follows from the definition of  $\gamma_n(\cdot|y)$  that  $\mu|_{\mathcal{F}_{\Lambda_n}}$  is not absolutely continuous with respect to  $\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}$ . Since the relative entropy is an increasing function of  $\mathcal{F}_{\Lambda_n}$  we have  $\mathcal{H}_{\Lambda_j}(\mu|\gamma_j(\cdot|y)) = +\infty$ ,  $\forall j \geq n$  and also  $\mathbf{h}^s(\mu) = -\infty$ , which proves the theorem in this case. Therefore we can assume  $\mathcal{H}_{\Lambda_n}(\mu|\gamma_n(\cdot|y)) < +\infty$ , for

all  $n \in \mathbb{N}$ . To lighten the notation we write  $d\mathbf{p} \equiv \prod_{i \in \mathbb{N}} dp$  to denote the product measure. Let  $\bar{f}$  a normalized potential cohomologous to  $f$  and  $\gamma_n(A|y) \equiv \mathcal{L}_{\bar{f}}^n(1_A)(\sigma^n(y))$ . If  $\mathcal{H}_{\Lambda_n}(\mu|\gamma_n(\cdot|y))$  is finite for all  $n \geq 1$  then

$$\begin{aligned}
\mathcal{H}_{\Lambda_n}(\mu|\gamma_n(\cdot|y)) &= \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} d\gamma_n(\cdot|y) \\
&= \mathcal{L}_{\bar{f}}^n \left( \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} \right) (\sigma^n(y)) \\
&= \int_{\Omega} \exp(S_n(\bar{f})) \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} \prod_{i \in \Lambda_n} dp \times \prod_{i \in \Lambda_n^c} d\delta_{y_i} \\
&= \int_{\Omega} \frac{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} \prod_{i \in \Lambda_n} dp \times \prod_{i \in \Lambda_n^c} d\delta_{y_i}. \\
&= \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} \prod_{i \in \Lambda_n} dp \times \prod_{i \in \Lambda_n^c} d\delta_{y_i},
\end{aligned}$$

where  $S_n(\bar{f}) \equiv \bar{f} + \bar{f} \circ \sigma + \dots + \bar{f} \circ \sigma^{n-1}$ . Since the above integrand is  $\mathcal{F}_{\Lambda_n}$ -measurable we get

$$\mathcal{H}_{\Lambda_n}(\mu|\gamma_n(\cdot|y)) = \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} d\mathbf{p}.$$

From the properties of the Radon-Nikodym derivative follows that

$$\begin{aligned}
\mathcal{H}_{\Lambda_n}(\mu|\gamma_n(\cdot|y)) &= \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \log \left( \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}} \frac{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \right) d\mathbf{p} \\
&\quad - \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \log \frac{d\gamma_n(\cdot|y)|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} d\mathbf{p} \\
&= \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} d\mathbf{p} - \int_{\Omega} S_n(\bar{f})(x_{\Lambda_n} y_{\Lambda_n^c}) d\mu(x) \\
&= -\mathcal{H}_{\Lambda_n}(\mu) - \int_{\Omega} S_n(\bar{f})(x_{\Lambda_n} y_{\Lambda_n^c}) d\mu(x),
\end{aligned}$$

where  $(x_{\Lambda} y_{\Lambda^c})_i = x_i$ , if  $i \in \Lambda$  and  $(x_{\Lambda} y_{\Lambda^c})_i = y_i$ , otherwise.

Note that

$$\int_{\Omega} \bar{f} d\mu = \lim_{n \rightarrow \infty} \frac{1}{n} \int_{\Omega} S_n(\bar{f})(x_{\Lambda_n} y_{\Lambda_n^c}) d\mu(x). \quad (3)$$

Indeed, for all  $n \geq 1$ , we have  $|S_n(\bar{f})(x_{\Lambda_n} y_{\Lambda_n^c}) - S_n(\bar{f})(x)| \leq \sum_{j=0}^n 2^{-\alpha j} \text{Hol}_{\alpha}(\bar{f})$ . From this observation and the  $\sigma$ -invariance of  $\mu$  the claim follows.

Recall that for any  $\mu \in \mathcal{M}_{\sigma}(\Omega)$  we have  $\mathcal{H}_{\Lambda_n}(\mu)/n \rightarrow \mathbf{h}^s(\mu)$ , when  $n \rightarrow \infty$ . This convergence together with (3) implies the existence of the following limit

$$\lim_{n \rightarrow \infty} \frac{\mathcal{H}_{\Lambda_n}(\mu|\gamma_n(\cdot|y))}{n} = -\mathbf{h}^s(\mu) - \int_{\Omega} \bar{f} d\mu = \log \lambda_f - \mathbf{h}^s(\mu) - \int_{\Omega} f d\mu,$$

where in the last equality we used that  $\mu \in \mathcal{M}_\sigma(\Omega)$  and the expression (2).

To finish the proof we only need to show that  $n^{-1} \mathcal{H}_{\Lambda_n}(\mu|\gamma_n(\cdot|y)) \rightarrow \mathfrak{h}(\mu|\nu)$ , when  $n \rightarrow \infty$ , for any choice of  $y \in \Omega$ .

$$\begin{aligned}
\mathcal{H}_{\Lambda_n}(\mu|\nu) &= \mathcal{H}_{\Lambda_n}(\mu|\nu\gamma_n(\cdot|y)) \\
&= \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d[\nu\gamma_n(\cdot|y)]|_{\mathcal{F}_{\Lambda_n}}} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d[\nu\gamma_n(\cdot|y)]|_{\mathcal{F}_{\Lambda_n}}} d[\nu\gamma_n(\cdot|y)] \\
&= \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \left( \frac{d[\nu\gamma_n(\cdot|y)]|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \right)^{-1} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d[\nu\gamma_n(\cdot|y)]|_{\mathcal{F}_{\Lambda_n}}} d[\nu\gamma_n(\cdot|y)] \\
&= \int_{\Omega} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d[\nu\gamma_n(\cdot|y)]|_{\mathcal{F}_{\Lambda_n}}} \frac{d[\gamma_n(\cdot|y)]|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} d\mu - \int_{\Omega} \log \frac{d[\gamma_n(\cdot|y)]|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} d\mu \\
&= \int_{\Omega} \log \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d[\gamma_n(\cdot|y)]|_{\mathcal{F}_{\Lambda_n}}} d\mu + \int_{\Omega} \log \frac{d[\gamma_n(\cdot|y)]|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \left( \frac{d[\nu\gamma_n(\cdot|y)]|_{\mathcal{F}_{\Lambda_n}}}{d\mathbf{p}|_{\mathcal{F}_{\Lambda_n}}} \right)^{-1} d\mu \\
&= \mathcal{H}_{\Lambda_n}(\mu|\gamma_n(\cdot|y)) \\
&\quad + \int_{\Omega} \left[ S_n(f)(x_{\Lambda_n} y_{\Lambda_n^c}) - \log \int_{\Omega} \exp(S_n(f)(x_{\Lambda_n} z_{\Lambda_n^c})) d\nu(z) \right] d\mu(x) \quad (4) \\
&= \mathcal{H}_{\Lambda_n}(\mu|\gamma_n(\cdot|y)) + o(n),
\end{aligned}$$

where the last expression follows from (3) together with the inequality  $|\log \int_{\Omega} \exp(\varphi) d\mu - \log \int_{\Omega} \exp(\psi) d\mu| \leq \|\varphi - \psi\|_{\infty}$ .  $\blacksquare$

**Corollary 1.** *For all  $f \in C^\alpha(\Omega)$  and  $\mu \in \mathcal{M}_\sigma(\Omega)$  we have  $\mathfrak{h}(\mu|\mu_{\bar{f}}) \geq 0$ .*

We now present the entropy considered in [LMMS15] and then we prove that it coincides with the specific entropy on the set of all Borel shift-invariant probability measures.

**Definition 2.** Given a Borel probability measure  $\mu$  on  $\Omega$ , we define its entropy as follows

$$\mathfrak{h}^v(\mu) \equiv \mathfrak{h}^{v,p}(\mu) \equiv \inf_{g \in C^\alpha(\Omega)} \left\{ - \int_{\Omega} g d\mu + \log \lambda_g \right\},$$

where  $\lambda_g$  is the maximal eigenvalue of  $\mathcal{L}_g$ .

**Theorem 3.** *For all  $\mu \in \mathcal{M}_\sigma(\Omega)$  we have  $\mathfrak{h}^s(\mu) = \mathfrak{h}^v(\mu)$ .*

*Proof.* Given  $\mu \in \mathcal{M}_\sigma(\Omega)$  it follows from Corollary 1 that  $\mathfrak{h}^s(\mu) \leq \log \lambda_g - \int_{\Omega} g d\mu$ , for all  $g \in C^\alpha(\Omega)$ . Therefore  $\mathfrak{h}^s(\mu) \leq \mathfrak{h}^v(\mu)$ .

The remainder of the proof is by contradiction. For the sake of simplicity, we will work on  $\mathcal{M}_s(\Omega)$ , the topological vector space of all finite Borel signed measures endowed

with the weak-\* topology, and with extensions  $\mathbf{h}^s, \mathbf{h}^v : \mathcal{M}_s(\Omega) \rightarrow [-\infty, 0]$  of  $\mathbf{h}^s$  and  $\mathbf{h}^v$ , respectively, given by

$$\mathbf{h}^s(\mu) = \begin{cases} \mathbf{h}^s(\mu), & \text{if } \mu \in \mathcal{M}_\sigma(\Omega) \\ -\infty, & \text{otherwise.} \end{cases} \quad \mathbf{h}^v(\mu) = \begin{cases} \mathbf{h}^v(\mu), & \text{if } \mu \in \mathcal{M}_\sigma(\Omega) \\ -\infty, & \text{otherwise.} \end{cases}$$

We split the proof in two separate cases.

**Case 1.** Suppose that there is  $\nu \in \mathcal{M}_s(\Omega)$  such that  $-\mathbf{h}^v(\nu) < -\mathbf{h}^s(\nu) < +\infty$ . Since the negative of the extension of the specific entropy  $-\mathbf{h}^s : \mathcal{M}_s(\Omega) \rightarrow [0, +\infty]$  is convex, lower semicontinuous and bounded from below, we have that its epigraph  $\text{epi}(-\mathbf{h}^s) \equiv \{(\mu, t) \in \mathcal{M}_\sigma(\Omega) \times \mathbb{R} : -\mathbf{h}^s(\mu) \leq t\}$  is a convex closed subset of  $\mathcal{M}_s(\Omega) \times \mathbb{R}$ . By assumption we have  $(\nu, -\mathbf{h}^v(\nu)) \notin \text{epi}(-\mathbf{h}^s)$ . Therefore, there are  $c \in \mathbb{R}$  and a continuous linear functional  $F : \mathcal{M}_s(\Omega) \times \mathbb{R} \rightarrow \mathbb{R}$ , such that for all  $(\mu, t) \in \text{epi}(-\mathbf{h}^s)$  we have  $F(\nu, -\mathbf{h}^v(\nu)) < c < F((\mu, t))$ . Since  $\mathcal{M}_s(\Omega)$  endowed with the weak-\* topology, the functional  $F$  can be represented as follows  $F((\mu, t)) = \int_\Omega \varphi d\mu + at$ , for some  $\varphi \in C(\Omega)$  and  $a \in \mathbb{R}$ . From the previous inequality follows that

$$\int_\Omega \varphi d\nu - a\mathbf{h}^v(\nu) < c < \int_\Omega \varphi d\mu + at, \quad \text{for all } (\mu, t) \in \text{epi}(-\mathbf{h}^s)$$

Since  $\mathcal{M}_\sigma(\Omega)$  is compact, and  $C^\alpha(\Omega)$  is dense in  $C(\Omega)$ , with respect to the uniform norm, up to small change in  $c$  we can assume that  $\varphi$  in above inequality is a Hölder continuous function. From this inequality it is easy to deduce that  $a > 0$ . Without loss of generality we can assume that  $a = 1$ . From Corollary 1, definition of  $\text{epi}(-\mathbf{h}^s)$ , the above inequality, and Theorem 3 of [LMMS15] we have

$$\log \lambda_{(-\varphi)} = \sup_{\mu \in \mathcal{M}_\sigma(\Omega)} \left\{ \int_\Omega (-\varphi) d\mu + \mathbf{h}^s(\mu) \right\} \leq -c < \int_\Omega (-\varphi) d\nu + \mathbf{h}^v(\nu) \leq \log \lambda_{(-\varphi)}$$

which is a contradiction. Therefore, for all  $\mu \in \mathcal{M}_\sigma(\Omega)$  such that  $\mathbf{h}^s(\mu) > -\infty$  we have  $\mathbf{h}^s(\nu) = \mathbf{h}^v(\nu)$ .

**Case 2.** Now we have to prove that for all  $\mu \in \mathcal{M}_\sigma(\Omega)$  such that  $-\mathbf{h}^s(\mu) = +\infty$ , we have  $-\mathbf{h}^v(\mu) = +\infty$ . The idea is to prove that we can reduce this to the previous case. Suppose that for some  $\nu \in \mathcal{M}_\sigma(\Omega)$  we have  $-\mathbf{h}^v(\nu) < \mathbf{h}^s(\nu) = +\infty$ . Since  $-\mathbf{h}^s$  is a convex and lower semicontinuous function from  $\mathcal{M}_s(\Omega)$  to  $[0, +\infty]$ , then it is pointwise supremum of a family  $\mathfrak{F}$  of continuous affine functions from  $\mathcal{M}_s(\Omega)$  to  $\mathbb{R}$ , see Proposition 3.1 of [ET76]. A generic member of  $\mathfrak{F}$  is a function of the type  $\mu \mapsto \xi(\mu) + C$ , where  $\xi \in \mathcal{M}_s(\Omega)^*$  and as observed before can be represented as  $\mu \mapsto \int_\Omega g d\mu + C$ , for some  $g \in C(\Omega)$ . Let  $\mathfrak{D}$  denote the family of all affine functions of the form  $\mathcal{M}_s(\Omega) \ni \mu \mapsto \int_\Omega g d\mu - \log \lambda_g$ , with  $g$  varying in  $C^\alpha(\Omega)$ . Since for all  $\mu \in \mathcal{M}_s(\Omega)$ , we have  $\mathbf{h}^s(\mu) \leq \mathbf{h}^v(\mu)$  then we can assume that  $\mathfrak{D} \subsetneq \mathfrak{F}$ . Given  $M > -\mathbf{h}^v(\nu)$  there is  $F \in \mathfrak{F} \setminus \mathfrak{D}$  such that  $F(\nu) > M$ . On the other hand, we have  $\mathbf{h}^s(\prod_{i \in \mathbb{N}} p) = 0$  so  $F(\prod_{i \in \mathbb{N}} p) \leq 0$ . Without loss of generality we can assume that the last inequality is strict, and for all  $\mu \in \mathcal{M}_s(\Omega)$ , we have  $F(\mu) = \int_\Omega \varphi d\mu + at$ , for some  $\varphi \in C^\alpha(\Omega)$  and  $a \in \mathbb{R}$ . Note that  $F$  is everywhere less than  $-\mathbf{h}^s$  and separates  $\text{epi}(-\mathbf{h}^s)$  from  $(\nu, -\mathbf{h}^v(\nu))$ . Therefore  $a \neq 0$  and the proof is finished by proceeding as in Case 1.  $\blacksquare$

**Theorem 4.** *If  $f \in C^\alpha(\Omega)$  then  $\mathfrak{h}(\mu|\mu_{\bar{f}}) = 0$  if and only if  $\mu \in \mathcal{G}^*(\bar{f})$ . In particular, the set of the equilibrium states for  $f$  is a singleton.*

*Proof.* If  $f \in C^\alpha(\Omega)$  then  $\mathcal{G}^*(\bar{f})$  is a singleton, see [LMMS15], and so  $\mathfrak{h}(\mu|\mu_{\bar{f}}) = 0$ .

Conversely, suppose that  $\mu \in \mathcal{M}_\sigma(\Omega)$  is such that  $\mathfrak{h}(\mu|\mu_{\bar{f}}) = 0$ . To prove that  $\mu \in \mathcal{G}^*(\bar{f})$  it is enough to show that for each fixed  $n_0 \in \mathbb{N}$  we have  $\mu\gamma_{n_0} = \mu$ , see reference [CL16] Definition 4 and remark b) below it, and Theorem 2. Since we are assuming that  $\mathfrak{h}(\mu|\mu_{\bar{f}}) = 0$ , and we know that  $\Lambda \mapsto \mathcal{H}_\Lambda(\mu|\mu_{\bar{f}})$  is a non-decreasing function (see [Geo11, p. 310]) it follows that  $\mathcal{H}_\Lambda(\mu|\mu_{\bar{f}}) < \infty$ , for any finite  $\Lambda \subset \mathbb{N}$ .

As a warm-up, let us first prove the theorem in the case  $\sup\{\mathcal{H}_{\Lambda_n}(\mu|\mu_{\bar{f}}) : n \in \mathbb{N}\} < +\infty$ . Under this assumption we have  $\mathcal{H}_{\Lambda_n}(\mu|\mu_{\bar{f}}) < +\infty$ , for all  $n \in \mathbb{N}$ , and so there exists the Radon-Nikodym derivative

$$\varphi_n \equiv \frac{d\mu|_{\mathcal{F}_{\Lambda_n}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_n}}}. \quad (5)$$

Recall that  $\varphi_n \geq 0$  and  $\mathcal{F}_{\Lambda_n}$ -measurable. It is a simple matter to check that  $(\varphi_n)_{n \in \mathbb{N}}$  is a martingale relative to  $\mu_{\bar{f}}$ . Indeed, since  $\mathcal{F}_{\Lambda_n}$  defines an increasing filtration, for every  $m > n$  we have  $\varphi_n = \mu_{\bar{f}}(\varphi_m|\mathcal{F}_{\Lambda_n})$   $\mu_{\bar{f}}$ -a.s. due to uniqueness of Radon-Nikodym derivative.

We will show that this martingale converges in  $L^1(\mu_{\bar{f}})$ -norm to some  $\mathcal{F}$ -measurable function  $\varphi \geq 0$ . To do this it is enough to prove that this sequence is uniformly  $\mu_{\bar{f}}$ -integrable. Indeed, for any  $K > 1$  we have

$$\begin{aligned} \int_{\Omega} \varphi_n 1_{\{\varphi_n \geq K\}} d\mu_{\bar{f}} &\leq \frac{1}{\log K} \int_{\Omega} [\varphi_n \log \varphi_n] 1_{\{\varphi_n \geq K\}} d\mu_{\bar{f}} \\ &< \frac{1}{\log K} \int_{\Omega} [1 + \varphi_n \log \varphi_n] 1_{\{\varphi_n \geq K\}} d\mu_{\bar{f}} \\ &\leq \frac{1}{\log K} (1 + \sup\{\mathcal{H}_{\Lambda_n}(\mu|\mu_{\bar{f}}) : n \in \mathbb{N}\}), \end{aligned}$$

where in the last inequality we used that  $1 + x \log x \geq 0$ , for all  $x \geq 0$ . From these estimates and the assumption that  $\sup\{\mathcal{H}_{\Lambda_n}(\mu|\mu_{\bar{f}}) : n \in \mathbb{N}\} < +\infty$ , we have that  $(\varphi_n)_{n \in \mathbb{N}}$  is a uniformly integrable martingale and thereby convergent in  $L^1(\mu_{\bar{f}})$ . Therefore there exists  $\varphi \in L^1(\mu_{\bar{f}})$  such that  $\lim_{n \rightarrow \infty} \int_{\Omega} |\varphi_n - \varphi| d\mu_{\bar{f}} = 0$ . Of course,  $\varphi = d\mu/d\mu_{\bar{f}}$ . Proposition 3 of [LMMS15] ensures that  $\mu_{\bar{f}}$  is ergodic. Since  $\mu$  is a shift invariant probability measure and  $\mu \ll \mu_{\bar{f}}$  it follows from a classical result in Ergodic Theory that  $\mu = \mu_{\bar{f}}$  and the theorem is proved in this case.

Now we go back to the general case. We shall prove that for any fixed  $n_0 \in \mathbb{N}$ , we have  $\mu\gamma_{n_0} = \mu$ . We split this proof in three steps.

**Step 1.** For each  $\delta > 0$  and each  $n_1 > n_0$ , there is  $n_2 \in \mathbb{N}$  such that  $n_0 < n_1 < n_2$  and

$$\mathcal{H}_{\Lambda_{n_2}}(\mu|\mu_{\bar{f}}) - \mathcal{H}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}(\mu|\mu_{\bar{f}}) \leq \delta, \quad (6)$$

where  $\mathcal{H}_{\Lambda_0}(\mu|\mu_{\bar{f}}) \equiv 0$ . Indeed, let  $m_1 \in \mathbb{N}$  be such that if  $n \geq \max\{n_1, m_1\}$ , then  $n^{-1}\mathcal{H}_{\Lambda_n}(\mu|\mu_{\bar{f}}) \leq \delta/n_1$ . By taking  $m_2 = \lceil \max\{n_1, m_1\}/n_1 \rceil$  we get

$$\frac{1}{m_2} \sum_{k=1}^{m_2} (\mathcal{H}_{\Lambda_{kn_1}}(\mu|\mu_{\bar{f}}) - \mathcal{H}_{\Lambda_{(k-1)n_1}}(\mu|\mu_{\bar{f}})) \leq \frac{1}{m_2} \mathcal{H}_{\Lambda_{m_2 n_1}}(\mu|\mu_{\bar{f}}) < \delta.$$

Therefore there is some  $n_2 > n_1$  such that  $\mathcal{H}_{\Lambda_{n_2}}(\mu|\mu_{\bar{f}}) - \mathcal{H}_{\Lambda_{n_2-n_1}}(\mu|\mu_{\bar{f}}) \leq \delta$ . Since  $\mu, \mu_f \in \mathcal{M}_\sigma(\Omega)$  it follows that  $\mathcal{H}_{\Lambda_{n_2-n_1}}(\mu|\mu_{\bar{f}}) = \mathcal{H}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}(\mu|\mu_{\bar{f}})$  and so the statement in step 1 is proved.

**Step 2.** Let  $\varphi_{n_2}$  be the function defined as in (5). Given  $\varepsilon > 0$  there exists  $\delta > 0$  such that

$$\int_{\Omega} \left| \varphi_{n_2} - \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}} \right| d\mu_{\bar{f}} < \varepsilon$$

whenever  $n_1 < n_2$  and  $\mathcal{H}_{\Lambda_{n_2}}(\mu|\mu_{\bar{f}}) - \mathcal{H}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}(\mu|\mu_{\bar{f}}) < \delta$ . In fact, if  $n_1$  and  $n_2$  are two positive integers satisfying  $n_1 < n_2$ . Then we have

$$\varphi_{n_2} = 0 \quad \mu_{\bar{f}} - a.s. \text{ on the set } \left\{ \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}} = 0 \right\}$$

because the last Radon-Nikodym derivative is equal to  $\mu_{\bar{f}}(\varphi_{n_2} | \mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}})$ . Consider the function  $\psi : [0, \infty) \rightarrow \mathbb{R}$ , given by  $\psi(x) = 1 - x + x \log x$ . For some  $0 < r < \infty$ , we have the following inequality  $|1 - x| \leq r\psi(x) + \varepsilon/2$ , for all  $x \geq 0$ . Therefore,

$$\begin{aligned} & \int_{\Omega} \left| \varphi_{n_2} - \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}} \right| d\mu_{\bar{f}} \\ &= \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}} \left| 1 - \varphi_{n_2} \left( \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}} \right)^{-1} \right| d\mu_{\bar{f}} \\ &\leq r \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}} \psi \left( \varphi_{n_2} \left( \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}} \right)^{-1} \right) d\mu_{\bar{f}} + \frac{\varepsilon}{2} \\ &= r \int_{\Omega} \varphi_{n_2} \log \left[ \varphi_{n_2} \left( \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}} \right)^{-1} \right] d\mu_{\bar{f}} + \frac{\varepsilon}{2} \\ &= r(\mathcal{H}_{\Lambda_{n_2}}(\mu|\mu_{\bar{f}}) - \mathcal{H}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}(\mu|\mu_{\bar{f}})) + \frac{\varepsilon}{2}. \end{aligned}$$

By taking  $\delta = \varepsilon/2r$  the step 2 statement is proved.

**Step 3.** To prove that  $\mu\gamma_{n_0} = \mu$  we fix a local function  $\phi$  (a bounded  $\mathcal{F}_{\Lambda_n}$ -measurable function for some  $n \in \mathbb{N}$ ) and  $\varepsilon > 0$ . Since  $\bar{f}$  is a Hölder potential it follows from Theorem 3 of [CL16] that the family of probability kernels  $(\gamma_n)_{n \geq 1}$  is quasilocal. Therefore there exists a local  $\mathcal{F}_{\Lambda_{n_0}^c}$ -measurable function  $\tilde{\phi}$  such that

$$\sup_{y \in \Omega} \left| \tilde{\phi}(y) - \int_{\Omega} \phi(x) d\gamma_{n_0}(x|y) \right| < \varepsilon.$$

Let  $n_1 > n_0$  be such that  $\phi$  is  $\mathcal{F}_{\Lambda_{n_1}}$ -measurable and  $\tilde{\phi}$  is  $\mathcal{F}_{\Lambda_{n_1} \setminus \Lambda_{n_0}}$ -measurable. Choose  $\delta$  in terms of  $\varepsilon$  as in Step 2, and define  $n_2$  in terms of  $n_1$  and  $\delta$  as in Step 1. Then we have

$$\begin{aligned}
\left| \int_{\Omega} \phi d\mu_{\gamma_{n_0}} - \int_{\Omega} \phi d\mu \right| &\leq \int_{\Omega} \left| \int_{\Omega} \phi(x) d\gamma_{n_0}(x|y) - \tilde{\phi}(y) \right| d\mu(y) \\
&\quad + \left| \int_{\Omega} \tilde{\phi} d\mu - \int_{\Omega} \tilde{\phi} \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}} d\mu_{\bar{f}} \right| \\
&\quad + \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}(y) \left| \int_{\Omega} \phi(x) d\gamma_{n_0}(x|y) - \tilde{\phi}(y) \right| d\mu_{\bar{f}}(y) \\
&\quad + \left| \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}(y) \left( \int_{\Omega} \phi(x) d\gamma_{n_0}(x|y) - \phi(y) \right) d\mu_{\bar{f}}(y) \right| \\
&\quad + \|\phi\|_{\infty} \int_{\Omega} \left| \varphi_{n_2} - \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}} \right| d\mu_{\bar{f}} \\
&\quad + \left| \int_{\Omega} \varphi_{n_2} \phi d\mu_{\bar{f}} - \int_{\Omega} \phi d\mu \right|.
\end{aligned}$$

Since  $\tilde{\phi}$  is  $\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}$ -measurable and  $\phi$  is  $\mathcal{F}_{\Lambda_{n_2}}$ -measurable, the second and last terms on rhs above are zero. Due to the choice of  $\tilde{\phi}$ , the first and third terms are each at most  $\varepsilon$ . The fifth term is not larger than  $\varepsilon \|\phi\|_{\infty}$  because of our choice of  $n_2$ . The fourth term is zero because of the DLR-equations. Indeed, since  $\mu_{\bar{f}} \in \mathcal{G}^*(\bar{f})$  and  $\bar{f}$  is Hölder follow from Theorem 2 of [CL16] that  $\mu_{\bar{f}}\gamma_{n_0} = \mu_{\bar{f}}$ , therefore

$$\begin{aligned}
&\left| \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}(y) \left( \int_{\Omega} \phi(x) d\gamma_{n_0}(x|y) - \phi(y) \right) d\mu_{\bar{f}}(y) \right| \\
&= \left| \int_{\Omega} \int_{\Omega} \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}(x) \phi(x) d\gamma_{n_0}(x|y) d\mu_{\bar{f}}(y) - \int_{\Omega} \phi \frac{d\mu|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}}{d\mu_{\bar{f}}|_{\mathcal{F}_{\Lambda_{n_2} \setminus \Lambda_{n_1}}}} d\mu_{\bar{f}} \right| = 0,
\end{aligned}$$

where in the first equality above we used that  $\gamma_{n_0}$  is a proper probability kernel and the DLR-equations.  $\blacksquare$

## 4 Concluding Remarks

In [LMMS15] the classical notion of *equilibrium states*, when  $\mathcal{A}$  is finite, is generalized for any compact metric alphabet. The authors fixed an a priori Borel probability measure  $p$  on  $\mathcal{A}$  and then defined equilibrium states for a Hölder potential  $f$  as being a  $\sigma$ -invariant probability measure solving the variational problem

$$\sup_{\mu \in \mathcal{M}_{\sigma}(\Omega)} \mathbf{h}^{\vee}(\mu) + \int_{\Omega} f d\mu.$$

They shown that any probability measure  $\mu$  satisfying  $\mathcal{L}_{\bar{f}}^* \mu = \mu$  is a solution to the variational problem. They also shown that there is a unique solution to  $\mathcal{L}_{\bar{f}}^* \mu = \mu$ , however they do not shown that any solution of the variational problem has to be of this form. We settled this question by combining the results of Theorems 1, 3 and 4. We remark that when  $\mathcal{A}$  is finite this result was first obtained in [GKLM18]. Our techniques also provided the uniqueness of the equilibrium states for potentials in Walters space, which is defined as follows.

**Definition 5** (Walters Space). A potential  $f : \Omega \rightarrow \mathbb{R}$  is said to be a Walters potential, notation  $W(\Omega, \sigma)$ , if the following condition is satisfied

$$\sup_{n \geq 1} \sup_{\mathbf{a} \in \mathcal{A}^n} |S_n(f)(\mathbf{a}x) - S_n(f)(\mathbf{a}y)| \rightarrow 0, \quad \text{when } d_\Omega(x, y) \rightarrow 0. \quad (7)$$

Theorem 1 ensures that

$$\lim_{n \rightarrow \infty} \frac{1}{n} \mathcal{H}_{\Lambda_n}(\mu|\nu) = \log \lambda_f - \int_{\Omega} f d\mu - \mathbf{h}^s(\mu),$$

where  $f$  is an  $\alpha$ -Hölder potential,  $\mu \in \mathcal{M}_\sigma(\Omega)$  and  $\nu \in \mathcal{G}^*(\bar{f})$ . The way we computed the above limit, the regularity properties of  $f$  are crucial. The main steps in this computation is the existence of  $\bar{f}$  (a normalized potential cohomologous to  $f$ ), the identity (3), and  $o(n)$  estimates for (4). The two former conditions are also verified for any potential in the Walters class and the existence of  $\bar{f}$  is proved in [CS16], for a general compact metric space alphabets. Therefore Theorem 1 can be generalized for such potentials. In the case of finite alphabet, the theorem can also be generalized to potentials in the Bowen class. Although it is not known whether the eigenfunctions associated to the maximal eigenvalue on this class is a continuous function its uniform upper and lower bounds, obtained in [Wal01], are enough for our argument. For a general compact metric space alphabet, as far as we know, the existence of the maximal eigenfunction has not been proved.

To obtain the uniqueness of the equilibrium states for a Walters potential  $f$ , we proceed as in the proof of Theorem 4, but using the equality  $\mathcal{G}^*(\bar{f}) = \mathcal{G}^{DLR}(\bar{f})$  and the uniqueness of  $\mathcal{G}^*(\bar{f})$  proved in [CL16].

Since Theorem 1 is based on the maximal spectral data of the Ruelle operator we have a natural way to extend the definition of  $\mathbf{h}(\mu|\nu)$ , when  $\nu$  is an equilibrium state of an arbitrary continuous potential  $f$ . For this extension instead of, using the maximal eigenvalue we use the spectral radius of  $\mathcal{L}_f$  acting on  $C(\Omega)$ . So we can define  $\mathbf{h}(\mu|\nu)$ , where  $\mu \in \mathcal{M}_\sigma(\Omega)$  and  $\nu$  an equilibrium state for some continuous potential  $f$ , by putting

$$\mathbf{h}(\mu|\nu) \equiv \log \rho(\mathcal{L}_f) - \int_{\Omega} f d\mu - \mathbf{h}^v(\mu).$$

Note that the above expression is similar to the one obtained in Theorem 1, but the specific entropy  $\mathbf{h}^s(\mu)$  is replaced by  $\mathbf{h}^v(\mu)$  and we can see that  $\mathbf{h}(\mu|\nu) \geq 0$ . On the other hand, we do not know, in general, whether  $n^{-1} \mathcal{H}_{\Lambda_n}(\mu|\gamma_{\Lambda_n}(\cdot|y))$  converges to  $\mathbf{h}(\mu|\nu)$ . Counterexamples, on the lattice  $\mathbb{Z}$ , given in Section A.5.2 of [vEFS93] (classical lattice systems) and in [MvE98] (quantum lattice systems) shows that this convergence can be a delicate issue.

## 5 Acknowledgements

The authors express many thanks to Aernout van Enter and Artur Lopes for their valuable comments and references. L. Cioletti and D. Aguiar are supported by CNPq and R. Ruviano is supported by FAP-DF and FEMAT.

## References

- [Bal00] V. Baladi. *Positive transfer operators and decay of correlations*, volume 16 of *Advanced Series in Nonlinear Dynamics*. World Scientific Publishing Co., Inc., River Edge, NJ, 2000.
- [CL16] L. Cioletti and A. O. Lopes. Ruelle operator for continuous potentials and DLR-Gibbs measures. *Preprint arXiv:1608.03881*, 2016.
- [CS16] L. Cioletti and E. A. Silva. Spectral properties of the Ruelle operator on the Walters class over compact spaces. *Nonlinearity*, 29(8):2253–2278, 2016.
- [ET76] I. Ekeland and R. Temam. *Convex analysis and variational problems*. North-Holland Publishing Co., Amsterdam-Oxford; American Elsevier Publishing Co., Inc., New York, 1976. Translated from the French, Studies in Mathematics and its Applications, Vol. 1.
- [Geo11] H-O. Georgii. *Gibbs measures and phase transitions*, volume 9 of *de Gruyter Studies in Mathematics*. Walter de Gruyter & Co., Berlin, second edition, 2011.
- [GKLM18] P. Giulietti, B. Kloeckner, A. Lopes, and D. Marcon. The calculus of thermodynamical formalism. *to appear in J. Eur. Math. Soc.*, 2018.
- [Isr79] R. B. Israel. *Convexity in the theory of lattice gases*. Princeton University Press, Princeton, N.J., 1979. Princeton Series in Physics, With an introduction by Arthur S. Wightman.
- [LMMS15] A. O. Lopes, J. K. Mengue, J. Mohr, and R. R. Souza. Entropy and variational principle for one-dimensional lattice systems with a general *a priori* probability: positive and zero temperature. *Ergodic Theory Dynam. Systems*, 35(6):1925–1961, 2015.
- [MvE98] H. Moriya and A. van Enter. On thermodynamic limits of entropy densities. *Lett. Math. Phys.*, 45(4):323–330, 1998.
- [PP90] W. Parry and M. Pollicott. Zeta functions and the periodic orbit structure of hyperbolic dynamics. *Astérisque*, (187-188):268, 1990.
- [Rue67] D. Ruelle. A variational formulation of equilibrium statistical mechanics and the Gibbs phase rule. *Comm. Math. Phys.*, 5:324–329, 1967.

- [Rue68] D. Ruelle. Statistical mechanics of a one-dimensional lattice gas. *Comm. Math. Phys.*, 9:267–278, 1968.
- [vEFS93] A. C. D. van Enter, R. Fernández, and A. D. Sokal. Regularity properties and pathologies of position-space renormalization-group transformations: scope and limitations of Gibbsian theory. *J. Statist. Phys.*, 72(5-6):879–1167, 1993.
- [Wal75] P. Walters. A variational principle for the pressure of continuous transformations. *Amer. J. Math.*, 97(4):937–971, 1975.
- [Wal01] P. Walters. Convergence of the Ruelle operator for a function satisfying Bowen’s condition. *Trans. Amer. Math. Soc.*, 353(1):327–347, 2001.