

ANALYSIS SEMINAR

Elliptic systems involving Schrödinger operators with vanishing potentials**Pedro Ubilla**

Universidad de Santiago de Chile

Date: 10/27/2022

Time: 02:00 pm

We prove the existence of a bounded positive solution of the following elliptic system involving Schrödinger operators

$$\begin{cases} -\Delta u + V_1(x)u = \lambda\rho_1(x)(u+1)^r(v+1)^p & \text{in } \mathbb{R}^N \\ -\Delta v + V_2(x)v = \mu\rho_2(x)(u+1)^q(v+1)^s & \text{in } \mathbb{R}^N, \\ u(x), v(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty \end{cases}$$

where $p, q, r, s \geq 0$, V_i is a nonnegative vanishing potential, and ρ_i has the property (H) introduced by Brezis and Kamin [1]. As in that celebrated work we will prove that for every $R > 0$ there is a solution (u_R, v_R) defined on the ball of radius R centered at the origin. Then, we will show that this sequence of solutions tends to a bounded solution of the previous system when R tends to infinity. Furthermore, by imposing some restrictions on the powers p, q, r, s without additional hypotheses of integrability on the weights ρ_i , we obtain a second solution using variational methods. In this context we consider two particular cases: a gradient system and a Hamiltonian system. This work is motivated in part by [2] and [3].

Joint work with Denilson Pereira and Juan Arratia.

References

- [1] H. Brezis and S. Kamin. *Sublinear elliptic equations in \mathbb{R}^N* . Manuscripta Math. 74 (1992), 87-106.
- [2] J. A. Cardoso, P. Cerda, D. S. Pereira and P. Ubilla. *Some Brezis-Kamin type problems and applications to Schrödinger Equation involving vanishing potentials*. Discrete Contin. Dyn. Syst, 2021, 41(6): 2947-2969.
- [3] M. Montenegro. *The construction of principal spectral curves for Lane-Emden systems and applications*. Ann. Sc. Norm. Super. Pisa Cl. Sci. 4 série, tome 29 (2000), 193-229.