

## On Mahler's $U_m$ -numbers

Ana Paula de Araújo Chaves (apchaves.math@gmail.com)

Universidade Federal de Goiás

**Abstract.** The genesis of transcendental number theory, took place in 1844 with Liouville's result on the "bad" approximation of algebraic numbers by rationals. More precisely, if  $\alpha$  is an algebraic number of degree  $n > 1$ , then there exists a positive constant  $C$ , such that  $|\alpha - p/q| > Cq^{-n}$ , for all  $p/q \in \mathbb{Q}^*$ . Using this remarkable fact, he was able to build a non-enumerable set of transcendental numbers called *Liouville numbers*. Since then, several classifications of transcendental numbers have been developed, one of them proposed by Kurt Mahler in 1932. He splitted the set of transcendental numbers on three disjoint sets:  $S$ -,  $T$ - and  $U$ -numbers. In a certain sense,  $U$ -numbers generalize the concept of Liouville numbers. Yet, the set of  $U$ -numbers can be splitted into  $U_m$ -numbers, that are numbers "rapidly" approximable by algebraic numbers of degree  $m$ .

On this lecture, the following result, made in cooperation with D. Marques, will be proved: Let  $\omega : \mathbb{N} \rightarrow \mathbb{N}$ , such that  $\omega_n \rightarrow \infty$ , as  $n \rightarrow \infty$ . Let  $\xi \in \mathbb{R}$  be a Liouville number, such that there exists an infinite sequence of rational numbers  $(p_n/q_n)_n$ , satisfying

$$\left| \xi - \frac{p_n}{q_n} \right| < H \left( \frac{p_n}{q_n} \right)^{-\omega_n},$$

where  $H(p_{n+1}/q_{n+1}) \leq H(p_n/q_n)^{O(\omega_n)}$ . Now, take  $\alpha_0, \dots, \alpha_l, \beta_0, \dots, \beta_r \in \overline{\mathbb{Q}}$ , with  $\beta_r = 1$  and  $\alpha_l \neq 0$ , such that  $[\mathbb{Q}(\alpha_0, \dots, \alpha_l, \beta_0, \dots, \beta_r) : \mathbb{Q}] = m$ . Then, for  $P(z), Q(z) \in \overline{\mathbb{Q}}[z]$ , given by  $P(z) = \alpha_0 + \alpha_1 z + \dots + \alpha_l z^l$  and  $Q(z) = \beta_0 + \beta_1 z + \dots + \beta_r z^r$ ,  $P(\xi)/Q(\xi)$  is a  $U_m$ -number.