

On selfmaps of complex Grassmann manifolds

Thaís Fernanda Mendes Monis*

Department of Mathematics

São Paulo State University (Unesp), Institute of Natural Sciences and Technology
Rio Claro-SP, Brazil

Abstract

Let $G(k, n)$ be the complex Grassmann manifold of k -planes in \mathbb{C}^{k+n} . A matter of interest is to know for which values of k and n the manifold $G(k, n)$ has the fixed point property. In this talk, we will discuss this problem, as well as related topics.

References

- [1] P. Chakraborty and P. Sankaran, Maps between certain complex Grassmann manifolds, *Topology Appl.*, 2014, 119–123.
- [2] H. Duan, Self-maps of the Grassmannian of complex structures, *Compositio Math.*, 2002, 159–175.
- [3] T. F. M. Monis, N. C. L. Penteado, S. T. Ura and P. Wong, A note on nontrivial intersection for selfmaps of complex Grassmann manifolds, *Bull. Belg. Math. Soc. Simon Stevin*, 2017, 665–672.
- [4] A. Taghavi, An alternative proof for the f.p.p. of $\mathbb{C}P^{2n}$, *Expo. Math.*, 2015, 105–107.

*Partially supported by CAPES, e-mail: thais.monis@unesp.br